






MHD simulation of cloud and clump formation triggered by the thermal instability

(and consequent massive star feedback)



Chris Wareing, J. Pittard, S.A.E.G. Falle, S. Van Loo
Astrophysics Group

‘Tracing the Flow’, Tuesday 3rd July 2018

Giant Molecular Clouds (GMCs)



UNIVERSITY OF LEEDS

Most stars are formed in GMC e.g. Rosette Molecular Cloud

Size $\simeq 35$ pc

Mass $\simeq 10^5 M_\odot$


Mean Density $\simeq 10^{-22}$ gm cm $^{-3}$

Temperature $\simeq 10$ K \Rightarrow sound speed $\simeq 0.2$ km s $^{-1}$

Alfvén speed $\simeq 2$ km s $^{-1}$ \Rightarrow magnetic pressure dominates

Velocity dispersion $\simeq 10$ km s $^{-1}$ - Supersonic and super Alfvénic

\Rightarrow Jeans Mass $\simeq 10^7 M_\odot$ (based on velocity dispersion)



Translucent Clumps



UNIVERSITY OF LEEDS

Rosette GMC not Homogeneous: *CO* maps show that it consists of $\simeq 70$ clumps with

Grey-scale: HI, contours CO, $1^\circ \equiv 28\text{pc}$

Sizes $\simeq 3.5 - 8.0 \text{ pc}$

Masses $\simeq 10^2 - 2 \cdot 10^3 \text{ M}_\odot$


Densities $10^{-21} \text{ gm cm}^{-3}$

Temperature $\simeq 10 \text{ K} \Rightarrow$ Sound speed $\simeq 0.2 \text{ km s}^{-1}$

Alfvén speed $\simeq 2 \text{ km s}^{-1} \Rightarrow$ magnetic pressure dominates (Crutcher 1999)

Velocity dispersion $\simeq 1 \text{ km s}^{-1}$ - Supersonic and sub Alfvénic

\Rightarrow Jeans Mass $3 \cdot 10^3 \text{ M}_\odot$ (based on velocity dispersion)



Blitz and Stark 1986

Physical model



UNIVERSITY OF LEEDS

Simplest approach with self-consistent physics for the formation of a molecular cloud and examine the results, before adding extra complexity

- 3D MHD
- Self-gravity
- Multi-phase ISM including thermal instability

In future, extra additions may be necessary:

- Shear and pressure waves, imitating galactic evolution
- Large-scale flows: SN shock, cloud collision
- “Turbulent” initial conditions applying randomised velocities

but if one can find a solution without recourse to extra complexity ...
lex parsimoniae / Occam’s razor

Thermal instability



UNIVERSITY OF LEEDS

Two stable phases exist in which heating balances cooling (Wolfire et al. 1995)

W – warm phase ($T > 5000\text{K}$, $\rho < 1$, $P/k < 5000$)

C – cold phase ($T < 160\text{K}$, $\rho > 10$, $P/k > 1600$)

U – unstable phase

In the unstable region, can form a length scale from cooling time and sound speed \sim a few pc.

Molecular cloud formation (10K) and stellar feedback (10^8K) requires multi-stage cooling:


$<10^4\text{K}$ Γ : Koyama & Inutsuka (2002), (2007 correction)

$10^4\text{K} < T < 10^8\text{K}$ Γ : CLOUDY 10.00 Gnat & Ferland (2012)

$>10^8\text{K}$ Γ : MEKAL - free-free bremsstrahlung.

Constant heating rate $\Gamma = 2 \times 10^{-26} \text{ erg s}^{-1}$ independent of ρ, T

\Rightarrow Establishes thermal equilibrium P and T by $\rho^2 \Lambda = \rho \Gamma$



Simple 3D Hydro condition



UNIVERSITY OF LEEDS

Spherical cloud, radius 50pc, density $n_H = 1.1$ - thermally unstable regime.

External medium density 0.1, pressure same as cloud. Self-gravity

Impose random 10% density perturbations
on finest initial AMR grid level (512^3)

Quiescent cloud $y=0$


Up to 10 levels of AMR (4096^3 : 0.037pc)

Mass: $1.7 \cdot 10^4 M_{\text{sun}}$

Sound crossing time: 6.458 Myrs

Free fall time: 44.92 Myrs

Cooling time: 1.642 Myrs





Simple 3D Hydro condition



UNIVERSITY OF LEEDS

First peak not due to gravity, second one is =>


$t=33.5$ Myrs



Detail at t=33.5 Myrs



UNIVERSITY OF LEEDS



Diameter ~ 5 pc, Mass $182 M_{\text{sun}}$, Max density 2214, Mean density 177, Max velocity 3.25 km s^{-1} (in frame of dense region), 0.6 km s^{-1} in dense gas.


Gravitationally bound, but not unstable (Bonnor-Ebert critical mass $\sim 471 M_{\text{sun}}$)

Enlarged 3D Hydro condition



UNIVERSITY OF LEEDS

Domain size doubled, cloud radius increased to 100pc ($r_{init} = 2.0$), initial maximum AMR resolution 1024^3 (finest level 0.29pc), Mass $1.35 \cdot 10^5 M_{sun}$



High density regions occur after
16.2 Myrs of diffuse cloud evolution
Increase resolution and simulate on...

- a further 28.5 Myrs
- resolution up to 0.039pc

Fellwalker (*how apt!*) clump identification
watershed algorithm (Berry 2015)


- 28 gravitationally isolated clumps
- size scale ~ 5 pc
- masses $50-300 M_{sun}$, $>80\%$ cold phase
- inward flow, dispersion $4-6 \text{ km s}^{-1}$
- ***unstable***

Will collapse to form clusters

Detail



UNIVERSITY OF LEEDS



Most massive clump: $354 M_{sun}$ (cold phase: $292 M_{sun}$), 5 pc diameter, max rho $1.5 \cdot 10^4$ (10^{-20} gm cm $^{-3}$), mean rho ~ 230 ($5 \cdot 10^{-22}$ gm cm $^{-3}$), dispersion 6.2 km s $^{-1}$.


3D MHD condition



UNIVERSITY OF LEEDS

Exactly the same as hydro, but with uniform field in the x-direction.

- Regular ($1.7 \cdot 10^4 M_{\text{sun}}$) and enlarged ($1.35 \cdot 10^5 M_{\text{sun}}$) clouds under consideration.
- Plasma β : 0.1 (strong field), 1.0 (plasma/magnetic pressure parity), 10.0 (weak field)



Magnetic seismology of Musca 'filament' indicates it is like this!


(Tritsis & Tassis 2018, Science, vol 360, Issue 6389, pp.635-638)

3D MHD condition – *in progress*



UNIVERSITY OF LEEDS

The plasma $\beta=10.0$ (weak field) case is looking very interesting





Gravitational collapse once the sheet has formed is dragging the field.

Field intensified in places from $0.3\mu\text{G}$ to $\sim 0.1\text{mG}$

$V_{\text{max}} \sim 3 \text{ km s}^{-1}$, $\mathcal{M}_{\text{max}} \sim 2.9$, $T \sim 10\text{K}$, $M \sim 150\text{M}_{\text{sun}}$

Density power spectrum $k^{-2.5}$



Beginning to show similarities?
=> Next step: re-simulate central section; sinks

Pattle, Ward-Thompson et al., 2017, ApJ, 846, id.122


Mechanical stellar wind feedback




UNIVERSITY OF LEEDS

Feedback simulations into these clouds have shown it's possible to clear a relatively small central cavity from a sheet-like parent molecular cloud.

What if the Rosette nebula...






...was formed by something like this:
($\beta=1$ cloud, $40M_{\text{sun}}$ feedback)



Simulating the Rosette Nebula



UNIVERSITY OF LEEDS



Evacuated hole

- Simulation: 10×7.5 pc
- Observations:
 - Celnik: $d \sim 13$ pc
 - IPHAS: $d \sim 10$ pc

Conclusions



UNIVERSITY OF LEEDS

Adopting only 3D hydrodynamics, thermal instability and self-gravity, it is possible to generate star-forming clumps from diffuse large-scale initial conditions.

With magnetic fields, sheets form, as recently inferred in the Musca cloud.

In the weak magnetic case, gravitational collapse intensifies field strength towards mG magnitudes and eventually will create double-horseshoe field structure.

A thin, extended molecular cloud in a magnetic field can host the Rosette Nebula.

But, how to create very low plasma β conditions? Pressure waves next!

Thank you for listening. Any comments or questions?

Thermal instability driven initial condition:

Wareing, Pittard, Falle & Van Loo, 2016, MNRAS, **459**, 1803

Magnetic feedback general case:

Wareing, Pittard & Falle, 2017, MNRAS, **465**, 2757

Hydrodynamic feedback general case:

Wareing, Pittard & Falle, 2017, MNRAS, **470**, 2283

Rosette special case:

Wareing, Pittard, Falle & Wright, 2018, MNRAS, **475**, 3598

Clumps formed by TI + gravity

Wareing, Pittard, Falle *in preparation*

The engine

Physical model



UNIVERSITY OF LEEDS

- Magnetohydrodynamic version of **MG** (*Morris Garages*) with self-gravity.
- Parallelised, upwind, conservative shock-capturing scheme.
- Adaptive mesh refinement uses a coarse base grid (4x4x4) with 7 (or more) levels of AMR to achieve a resolution up to 512^3 (*the Honda bit?*).
 - Why the wide range? Efficient computation of self-gravity.
- Realistic heating and cooling methods
 - Of key importance as it is the balance of these that establishes the initial condition and defines the consequent evolution.
- Three field strengths considered, with $\underline{B} = B_o \hat{I}_x$ $\beta = \frac{\rho k_B T}{B^2/2\mu_0}$ $\frac{\text{thermal pressure}}{\text{magnetic pressure}}$
 - The hydrodynamic case: $\beta = \infty$
 - Pressure equivalence: $\beta = 1$ - inferred to be the commonest in reality.
 - Magnetically dominated regime: $\beta = 0.1$

Aside: EPSRC and Innovate UK
research proposals to apply MG in
industry: cryogenic machining.



HONDA
The Power of Dreams
TYPE R